



**University of
Zurich** ^{UZH}



TROUBLESHOOTING AND ENGINEERING OF ANTIBODY CONSTRUCTS - PART II


Jonas V. Schaefer, PhD

Head of High-Throughput Laboratory
Department of Biochemistry, University of Zurich

www.bioc.uzh.ch/plueckthun



Antibody therapeutics vs. engineering




Total of 165 anti-cancer antibodies currently in clinical studies:

- 84 unmodified IgG (51%)
- 25 ADC (15%)
- 10 bispecific (6%)
- 17 engineered (10%)
- 16 fragments (10%)

➔ while most antibodies on the market / in R&D are full-length IgGs, most of the antibody engineering is performed using small fragments

Full-length IgG engineering

Optimization antigen binding
(affinity, humanization, decreased elimination)



➔ variable domains mainly used
for individual optimization


➔ most "transferable" engineering
is focused on F_C region




Why not just one "perfect" framework?

seven V_H germline families with different biophysical properties


➔ variability in subfamilies increases
binding diversity



Haptent




Peptides



Proteins


Engineering of unstable V_H6 domain

comparison of the human consensus V_H domains (germinal)




improved biophysical properties of scFv fragments expressed in *E. coli*:

- increased stability: $\Delta\Delta G_{N-U} = 20.9$ kJ/mol
- 4-fold increase in expression levels





Are previous findings transferable?





Model antibodies




	IgG 6B3	IgG 2C2
heavy chain (HC)	V _H 6	V _H 6
antigen	protein	peptide
light chain (LC)	V _λ 3 (lambda)	V _κ 3 (kappa)

➔ chosen model IgGs differ in

- Fab stability: rather unstable (6B3) vs. extremely stable (2C2)
- pI: 6.9 (6B3) vs. 8.7 (2C2)
- antigen: protein vs. peptide




Comparison of expression levels



➔ eukaryotic chaperons and quality control systems equalize the expression yield between WT and stabilized V_{H6}



➔ prokaryotic expression of IgGs indicates increased periplasmatic levels of the M variants





Analytical challenge: Multidomains

- ➔ IgGs consist of six individual domains (each in duplicates), all having similar folds




- ➔ with most experimental setups, only **overall average** of biophysical features will be analyzed



Biophysical analyses (methodology)

Circular Dichroism (CD)




(2^{ry} structure composition)

thermal denaturation

(aggregation analysis)



Intrinsic Tryptophan Fluorescence (ITF)



thermal denaturation

chemical denaturation

Differential scanning calorimetry / fluorimetry



analysis of individual domains




Circular Dichroism (CD)

Lambert-Beer derivative: $\Delta A = A_L - A_R = \epsilon_L \times l \times C - \epsilon_R \times l \times C = \Delta\epsilon \times l \times C$

ellipticity: $\theta = \frac{2.303 (A_L - A_R)}{4l}$

MRE: $[\theta] = \frac{\theta \times 100 \times M}{C \times l \times n}$


amide chromophore of peptide bond has 2 electronic transitions of low energy:
 $n \rightarrow \pi^*$ (signals at 222 nm and 215 nm) and $\pi \rightarrow \pi^*$ (signals at 208 nm and 198 nm)





- ➡ at ~ 208 nm intensity due to β-sheets is essentially zero
- ➡ measuring ellipticity at 208 nm monitors **changes in structure** (negative shift caused by random coil formation)



CD: real examples



➔ unfolding detectable, however sheaded by aggregation





Trp fluorescence is very sensitive to local conformation and environment

Quantum yields:

Phe - 0.02

Tyr - 0.13

Trp - 0.12

IgG 2C2: 24 Trp per IgG

IgG 6B3: 26 Trp per IgG

IgG 6B3

Domain	# of Trp	% of all Trp
V _H	5	38.5
CH ₁	1	7.7
CH ₂	2	15.4
CH ₃	2	15.4
V _L	1	7.7
CL	2	15.4

IgG 2C2

Domain	# of Trp	% of all Trp
V _H	5	41.7
CH ₁	1	8.3
CH ₂	2	16.7
CH ₃	2	16.7
V _L	1	8.3
CL	1	8.3




➔ majority of Trp residues are located within V_H domain



Trp fluorescence is very sensitive to local conformation and environment

wavelength maximum shifts upon heating due to changes of polarity in vicinity of Trp (**red-shift** of Trp emission spectrum)

red shift can be monitored by ratio of intensities at 330 and 350 nm





benefit over other methods:

- aggregation doesn't cover unfolding reaction
- can easily be performed in 96well format





ITF: real examples

Temperature-unfolding




GdnHCl-unfolding






Real-time GdnHCl denaturation


HEK IgG 6B3 WT




HEK IgG 6B3 M



HEK IgG 2C2 WT




HEK IgG 2C2 M





Differential Scanning Fluorimetry (DSF)



melting temperature detected by increased fluorescence of dye with **affinity for hydrophobic parts of the protein**



Sypro-Orange (Molecular Probes)





in aqueous solution: **quenched fluorescence;**
highly fluorescent in non-polar environment

relatively **high excitation wavelength**
decreases likelihood of small molecules
interfering with optical properties of dye,
causing quenching of fluorescence intensity






DSF: real examples



Power-compensation DSC (not Heat-flux DSC)



continuously self-adjustment
of heating power for keeping
sample and reference at same
temperature

difference of required power
[J/sec] divided by the scan
rate [°C/sec] leads to heat
capacity [J/°C]

Integration of heat capacity vs. temperature yields the enthalpy (ΔH)




$$\Delta H = \int_{T_1}^{T_2} C_p dT$$

$$\Delta G = \Delta H - T \cdot \Delta S$$

(Gibbs Free Energy equation)



DSC: real examples



➡ DSC is only setup detecting small differences of very stable transition





VP-DSC vs. VP-Capillary DSC



plate storage

autosampler









VP-DSC vs. VP-Capillary DSC

	VP-DSC	VP-Capillary DSC
analyzed volume	510 μl	130 μl
sample volume	1'200 μl	400 μl
scan rates	0.5 - 1.5 $^{\circ}\text{C}/\text{min}$	0.16 - 4 $^{\circ}\text{C}/\text{min}$
sample cell	coin shaped	capillary
samples	1	up to 288
measuring time	1 day	4 hrs
cleaning	manual	automatic

major advances: sensitivity, throughput, reproducibility, stability and ease of use
(smaller sample requirements)



VP-DSC vs. VP-Capillary DSC





Convection at aggregation

protein aggregation: heat signal detected by DSC is sum of both **endothermic unfolding** and **exothermic aggregation**

convection appears



once sample aggregates, interference and baseline drop

molecules are located in small confined space

very little convection





due to small diameter of capillaries

molecules are separated with enough space (aggregation delayed)

➔ signals derived from Capillary-DSC are **less sensitive to aggregation**



Comparison DSF vs. DSC



compared to DSC, DSF lacks "resolution" of individual domains, however is much faster (2-3 hrs vs. 48-72 hrs), can be performed in parallel and requires much less protein (20 μg vs. ~ 1 mg)



Stabilizing effects of V_H6 mutations

		ITF	GdnHCl	DSF	DSC
IgG 2C2	WT	70.4°C*	2.5 M	n.d.	86.0°C
	M	71.8°C*	3.8 M	n.d.	87.8°C
		Δ = 1.4°C	1.3 M	-	1.8°C
<hr/>					
IgG 6B3	WT	67.6°C	2.0 M	74.5°C	72.1°C
	M	70.8°C	2.6 M	77.0°C	74.3°C
		Δ = 3.2°C	0.6 M	2.5°C	2.2°C
<hr/>					
Fab 6B3	WT	69.7°C	2.0 M	76.5°C	72.6°C
	M	74.2°C	2.6 M	80.0°C	76.6°C
		Δ = 4.5°C	0.6 M	3.5°C	4.0°C

* - determined in presence of 1 M GdnHCl

n.d. - not determined




**IgG stability
analyses**



**IgG expression
systems**




Mammalian cell culture



stable HEK293 (Flp-In)

CMV promoters (constitutive)



Yeast *Pichia pastoris*



stable SMD1163 (his4 pep4 prb1)

GAP promoters (constitutive)

Difference in aggregation susceptibility



- ➔ *Pichia*-derived glycans reduce aggregation tendency
- ➔ peptide remaining from yeast signal sequence decreases aggregation susceptibility of HEK-IgG upon N-terminal addition



Expression of full-length IgGs in methylotrophic yeast *Pichia pastoris*

➔ advantages of expression system:

- disulfide bond formation / isomerization
- posttranslational modification (glycosylation)
- very high cell densities
- high expression levels (up to 30%)

➔ different promoters available:



- MeOH-inducible AOX1 (alcohol oxidase 1)
- constitutive GAP (glyceraldehyde-3-phosphate dehydrogenase)

➔ only low-level secretion of endogenous proteins, being advantageous for protein purification and downstream processing

➔ > 50 reports describing antibody expression
(mainly scFvs, several Fabs, only handful full-length IgG)


major difference in expression systems: glycosylation

- ➔ yeast system processes same sugar precursor differently (in Golgi complex), resulting in a **different glycan**






N-linked glycosylation






N-linked glycan processing

HEK293 cells




$\text{Gal}(\text{GlcNAc})_2(\text{Man})_3(\text{GlcNAc})_2\text{Fuc}$

Pichia pastoris



$(\text{Man})_{9-10-18}(\text{GlcNAc})_2$







➔ *Pichia* glycan cause difficulties interacting with Fcγ receptors (FcγR) important for effector functions



Influence of glycosylation on stability

- ➔ *Pichia* produced IgGs have decreased C_H2 stability, compared to mammalian expression
- ➔ different C_H2 stabilities are caused by different glycan moieties






**IgG stability
analyses**



**IgG
homogeneity**


Electrophoretic analyses of IgGs





➔ non-reducing SDS-PAGE reveals inhomogeneity of WT, but not of M variants

➔ banding pattern is not caused by:

- glycosylation
- proteolysis
- charge heterogeneity




Disulfide bond scrambling in IgG2



➔ Disulfide shuffling as source of multiple bands?

Analysis of IgG variants on "non-conventional" 2D-SDS-PAGE

HEK IgG 2C2 WT





- ➔ distinct bands of 1st dimension resolve again in the 2nd dimension
- ➔ multiple bands are **not caused by disulfide heterogeneity / shuffling** (confirmed by MS analyses and determination of unpaired cysteines)



Stability probed by dye binding

Analysis by capillary electrophoresis (performed in microfluidic chip)






➡ M variant seems more densely packed (less SDS-micelles can bind)




Stability probed by partial reduction

Partial reduction of IgG by hydrophilic TCEP



➡ TCEP treatment reduces inter-molecular disulfide bond only in WT IgGs


➡ labeling of free Cys with fluorescent 5-IAF confirms improved structural integrity / compactness





Conclusions


mutations affect structural integrity and homogeneity





Conclusions

- variable domain mutations: effects on expression level
 - strong influence in *E. coli*
 - moderate influence in *Pichia pastoris*
 - no influence in HEK293
- mutations influence the biophysical properties of the IgG: thermal and denaturant-induced unfolding
- increased stability independent of the expression system used
- **transferability of improvements implemented in smaller fragments onto full-length IgG**





Acknowledgements

Dept. of Biochemistry, UZH

Andreas Plückthun

Birgit Dreier

Annemarie Honegger

Peter Lindner

all present and former lab members

Academic partners

Ilian Jelezarov (UZH)

Paolo Cinelli (UZH)

Functional Genomics Center (UZH)

Shaikh Rafeek (ZHAW)

Manfred Heller (University of Bern)

Yuguang Zhao (Wellcome Trust, Oxford)

Margaret Jones (Wellcome Trust, Oxford)



general / financial support



FCI
FONDS DER
CHEMISCHEN
INDUSTRIE



Industrial partners

Peter Gimeson (GE Healthcare)



Daniel Weinfurtner (MorphoSys)



Thomas Müller-Späth (ChromaCon)



ChromaCon

Stefan Duhr (NanoTemper)





Questions & Answers



Jonas V. Schaefer

jonas.schaefer@uzh.ch



Schaefer and Plückthun (2012) *Protein Eng. Sel. Des.* 25(10):485-506

Schaefer and Plückthun (2012) *J Mol Biol.* 417(4):309-35